



## Isothermal Adsorption Modelling of *Pupuru* Flour

 **ADESHINA FADEYIBI<sup>1</sup>\*** AND **JANET OLWATOYIN ALABA<sup>2</sup>**

<sup>1</sup>Department of Food and Agricultural Engineering, Kwara State University, Malete, Nigeria

<sup>2</sup>Department of Food Science and Technology, Wesley University of Science and Technology, Ondo state, Nigeria

### ARTICLE INFO

#### Article history:

Received: April 14, 2020

Accepted: July 31, 2020

#### Keywords:

Adsorption moisture sorption,  
Equilibrium moisture content,  
Monolayer,  
*Pupuru*

### ABSTRACT

The amount of moisture adsorbed by stored food products usually varies with temperature and relative humidity at a constant pressure. This research was carried out to determine the sorption isotherm behaviour of *pupuru* flour in the temperature and water activity ( $a_w$ ) ranges of 10–50 °C and 0.10–0.93 using gravimetric method. The Guggenheim-Anderson-De Boer (GAB), Brunauer-Emmett-Teller (BET), Oswin and Smith sorption equations, formulated for the sorption characterization of starchy foods, were used to constitute the models. The results show that the sorption isotherm curve is similar to a Type II classification typical of starchy foods. Also, the sorption parameters ( $M_o$ , C and K) were generally higher for the products stored at 10 °C and 0.10  $a_w$ . The parameter K decreases with an increase in the temperature for the GAB and the Smith models, but remains inconsistent for the Oswin model. The  $M_o$  increases with an increase in the temperature for the GAB and Smith models. The GAB, Oswin and Smith models were generally adequate and valid with an  $MSE < 10\%$  and a high degree of fitness ( $R^2$  and  $R^2_{adj} > 90\%$ ). This indicates their ability to predict the sorption isotherm characteristics of the *pupuru* flour within its range of the  $a_w$  and the storage temperature.

## Introduction

*Pupuru* is a fermented, dewatered, milled and toasted flour produced from cassava. It originates from the Ilaje people of the riverine area of Ondo State, South-west Nigeria (Shittu et al., 2001). The traditional processing of *pupuru* involves steeping the peeled cassava roots in water for four days after which the wet pulp is dried by spreading it on a raised platform, constructed using a raffia material locally known as *Aka*. Based on the processing method of *pupuru*, it is different from other cassava fermented products like *garri*, *lafun*, *fufu* and *akpu* (Osundahunsi, 2005). The fermented cassava mash can be mould into balls and smoke-dried. This is believed to impact some characteristics of flavour and aroma to this product (Shittu et al., 2003; Daramola et al., 2010). According to Famurewa et al. (2013), producing *pupuru* flour by toasting method can help solve the problem associated with making wet *pupuru* before smoking. A common problem is the deterioration of the product, and this is responsible for

its occasional low market value and acceptability. The deterioration of the product may be associated with the activities of the spoilage organisms such as bacteria and fungi (Ilouno et al., 2016; Adebowale et al., 2017). An understanding of the sorption isotherm characteristics of the product is therefore critical to prevent the deterioration, especially before processing and storage (Oluwamukomi et al., 2008). The water sorption isotherm of foods is therefore of great importance.

A moisture sorption isotherm describes the thermodynamic relationship between water activity and the equilibrium moisture content of a food product at constant temperature and pressure (Erbaş et al., 2016). This relationship can be nonlinear or linearly constituted by two or more partial regression coefficients. Several mathematical models have been proposed for describing sorption isotherms of food products. Some of them were developed with a theoretical basis to describe adsorption mechanisms (Erbaş et al., 2016; Rockland and Stewart, 2013), whereas the others are just empirical or a simplification

\*Corresponding author E-mail: [adeshina.fadeyibi@kwasu.edu.ng](mailto:adeshina.fadeyibi@kwasu.edu.ng)

of more elaborate models. In some ranges of water activity, sorption isotherms can be approximated to linear equations (Yan et al., 2008). There are some semi-empirical equations with two or three fitting parameters which describe the moisture sorption isotherms of foods. The fundamental equations have been reported (Erbaş et al., 2016; Aguirre-Loredo et al., 2017; Mitrevski et al., 2017; Tejada-Ortigoza et al., 2017), but their application for predicting the adsorption and desorption characteristics depends on the nature of the food, water activity, equilibrium moisture content and the temperature of storage.

Guggenheim-Anderson-De Boer (GAB) and Brunauer-Emmett-Teller (BET) models have been applied most successfully to describe the equilibrium isotherm behaviour of many food products (Tejada-Ortigoza et al., 2017). In fact, these models have been recommended by the European Project Group COST 90 on the Physical Properties of Foods as the fundamental equation for the characterization of water sorption of food materials (Samapundo et al., 2007). The adsorption and desorption isotherms of maize flour and yellow dent corn have been investigated by Oyelade et al. (2008) and Samapundo et al. (2007). The authors recommended the modified GAB model for predicting the safe moisture content for the storage of the products. Similarly, the effect of hydrophobic modifications on the adsorption isotherms of cassava starch has been reported (Cova et al., 2010). The authors reported that the water adsorption process was slower for modified starches. Although the storage stability of *pupuru* flour has been studied at room temperature (Famurewa et al., 2012), this study is limited, because the authors did not consider the storage effect at higher temperatures. Despite the wider application of the sorption isotherm models in predicting the sorption behaviour of stored food products, there are no reported research on their application to predict the sorption characteristics of *pupuru* flour. It is therefore imperative to determine the optimal storage conditions that would best maintain the quality attributes of the *pupuru* flour during storage and processing. Hence, the objective of this study was to determine sorption isotherm of *pupuru* at three different temperatures and water activities, and to validate sorption characteristics of the product using four different sorption equations for the starchy foods.

## Materials and methods

### Preparation of Pupuru

The preparation of the *Pupuru* flour was done as described by Famurewa et al. (2012). This involves peeling of the cassava tubers and washing them in clean water for 4 days at ambient temperature until

they were softened. The water was changed every 24 hours to reduce the odour and allow the leaching of the antinutrient content out. After the fourth day, the water was drained and the fermented mash was pressed with a hydraulic press. The hard fibers were manually removed and the wet cassava meal sieved with a mesh to remove shaft. The meal was toasted in a wide aluminum pot for 20 min with a constant stirring to prevent burning and formation of lumps. Thereafter, the product was cooled and milled into flour. The sample of the *pupuru* flour was then packaged in HDPE film and kept in refrigerated storage for subsequent experimentation.

### Determination of Equilibrium Moisture Content

The equilibrium moisture content of the *pupuru* flour was determined according to the procedures reported by Singh and Talukdar (2020) for static gravimetric analysis. The method involves the application of a saturated salt solution on the flour sample to establish a constant value of the water activity ( $a_w$ ), until equilibrium is achieved between the atmosphere and the food. For the adsorption process, the *pupuru* flour samples were dehydrated in a hot air oven to bone dry state. Saturated solutions of LiCl, MgCl<sub>2</sub>, Mg(NO<sub>3</sub>)<sub>2</sub>, Na<sub>2</sub>NO<sub>3</sub>, NaCl and KNO<sub>3</sub> salts were prepared and enclosed in a desiccator. The  $a_w$  of salt solutions were found to vary between 0.113 to 0.932 at three different temperatures, namely 10 °C, 30 °C and 50 °C. The sample of the *pupuru* flour (3 g) was placed into containers with the corresponding saturated salt solutions. Toluene was placed into the containers that had  $a_w$  values higher than 0.65 to prevent microbial growth. The desiccators were maintained at  $a_w$  values of 0.1 to 0.93. Each of the samples was weighed using a digital balance until a constant weight was achieved, thus indicating equilibrium. The bone-dry mass was determined by oven drying method (AOAC, 2005). The procedure was repeated three times, and the average values and standard deviation of the equilibrium moisture content was computed at different temperatures and water activities of the *pupuru* flour samples.

### Determination of Sorption Parameters of Pupuru Flour

#### GAB sorption model

The results of the moisture adsorption were fitted to the GAB equation (Fadeyibi et al., 2012; Bello et al., 2018). The GAB equation can be expressed as shown in Equation (1).

$$M_e = \frac{CkM_o a_w}{(1-ka_w)(1-ka_w+Cka_w)} M_o \quad (1)$$

The three parameters of the GAB model (c, k and  $M_o$ ) were obtained from the second order polynomial expression, which was solved by multi-linear regression analysis to obtain  $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $R^2$  and E% (Xiong, 2002) according to equations (2) – (5).

$$\frac{a_w}{M_e} = \alpha a_w^2 + \beta a_w + \gamma \quad (2)$$

where

$$\alpha = \frac{k}{M_o} \left( \frac{1}{c-1} \right) \quad (3)$$

$$\beta = \frac{1}{M_o} \left( 1 - \frac{2}{c} \right) \quad (4)$$

$$\gamma = \frac{1}{M_o} Kc \quad (5)$$

The values of the GAB model parameters  $M_o$ , C and K, for each temperature, were therefore determined using equations (6) to (8).

$$M_o = \frac{1}{\sqrt{\beta^2 - 4\alpha\gamma}} \quad (6)$$

$$C = \frac{2\sqrt{\beta^2 - 4\alpha\gamma}}{-\beta + \sqrt{\beta^2 - 4\alpha\gamma}} \quad (7)$$

$$K = \frac{-\beta \pm \sqrt{\beta^2 - 4\alpha\gamma}}{2\alpha\gamma} \quad (8)$$

#### BET sorption model

The monolayer moisture content was also determined using the BET sorption model as expressed in equation (9) (Bello et al., 2018). The values of the equilibrium moisture content were plotted against water activity ( $a_w$ ) to obtain sorption isotherm curves of the *pupuru* flour for different storage temperatures. The BET curves were obtained by plotting the graph  $a_w/(1-a_w)M_e$  against  $a_w$ . The slope and intercept were determined from the BET curves using linear regression analysis. Monolayer moisture content ( $M_o$ ) and the constant (C) were calculated from the analysis (Labuza and Altunakar, 2020).

$$M_e = \frac{M_o C a_w}{(1-a_w)(1+(C-1)a_w)} \quad (9)$$

$$\frac{a_w}{(1-a_w)M_e} = \frac{1}{CM_o} + \frac{a_w(C-1)}{CM_o}$$

$$\frac{a_w}{(1-a_w)M_e} = \alpha a_w + \beta$$

where

$$\beta = \frac{1}{CM_o}$$

$$\alpha = \frac{(C-1)}{CM_o}$$

$M_o$  = monolayer moisture content,

C = energy constant,

$M_e$  = equilibrium moisture content (moisture content adsorbed per dry weight) in gH<sub>2</sub>O/g solids.

#### Oswin sorption model

The sorption data obtained were also fitted into the Oswin model as expressed in Equation (10). This model is essentially useful in describing the sorption isotherm behaviour of fat free and starchy foods. Thus, the model was used to determine the monolayer moisture content of *pupuru* flour.

$$M_e = C \left( \frac{a_w}{1-a_w} \right)^n \quad (10)$$

$$\ln M_e = \ln C + n \ln \left( \frac{a_w}{1-a_w} \right)$$

$$\ln M_e = \alpha \ln \left( \frac{a_w}{1-a_w} \right) + \beta$$

Thus, a plot of  $\ln M_e$  versus  $\ln (a_w/1-a_w)$  gives a straight line with slope  $n = k = \alpha$  and intercept  $\ln C = M_o = \beta$ .

#### Smith sorption model

The Smith sorption equation was used to predict the adsorption isotherm characteristics of the *pupuru* flour for different temperature conditions. The equation was used because it aids the understanding of the sorption behaviour of starchy foods which consist of a single and multiple layers of condensed water molecules. In this study, we computed the Smith sorption model parameters using Equation (11).

$$M_e = C_1 + C_2 \ln(1 - a_w) \quad (11)$$

Putting  $C_1 = M_o$ , and  $C_2 = k$ , we have

$$M_e = M_o + k \ln(1 - a_w)$$

where

$C_1$  = quantity of water in the first sorbed fraction,

$C_2$  = quantity of water in the multilayer moisture fraction.

### Model Verification and Calibration

The validity of the GAB, BET, Oswin and Smith models for predicting the equilibrium and monolayer moisture contents of the *pupuru* flour for the different temperature conditions was tested. This was done by comparing the values with the corresponding values from an independent experiment (Ajisegiri et al., 2007). The fitness of the models was tested using the mean relative percentage error (MRE, %), and the coefficients of determination ( $R^2$ ) and  $R_{adj}^2$  as expressed in Equations (12) to (14), respectively.

$$MRE = \frac{100}{k} \sum_{i=1}^N \left( \frac{M_{ei} - M_{ci}}{M_{ei}} \right)^2 \quad (12)$$

$$R^2 = \sum \frac{(M_{pred} - M_{avg})^2}{(M_{avg} - M_{av})^2} \quad (13)$$

$$R^2_{adj} = 1 - \frac{(1-R^2)(k-1)}{k-p-1} \quad (14)$$

where

$M_{ei}$  = moisture content (% db)

$M_{ci}$  = calculated moisture content (% db)

$k$  = number of samples

$p$  = number of independent variables

$M_{pred}$  = model predicted moisture content (% db)

$M_{avg}$  = average moisture content (% db)

### Results and discussion


#### Sorption Isotherm Characteristics of Pupuru

The adsorption isotherm of *pupuru* samples are presented in Fig. 1. There was an increase in the equilibrium moisture content with an increase in the  $a_w$ . The sorption isotherm curve showed the typical sigmoid


shape confirming type II classification which is the characteristic of the most biological material. This absorbs relatively small amount of water at lower activities and large amounts at high relative humidity. This agrees with the works of Kumar et al. (2005) for products rich in starch content. The moisture sorption characteristics of a product have been shown to be influenced by its composition, processing treatment, temperature and relative humidity (Ojuko and Olapade, 2018). The more the quantity of the adsorbed moisture, the higher the value of water activity.

#### Sorption Parameters of Pupuru Flour


The adsorption isotherm of the *pupuru* flour stored at different temperatures based on the GAB, BET, Oswin and Smith models are shown in Figs. 2–5, respectively. Quadratic relationships were established between the equilibrium moisture content and the  $a_w$  at different storage temperatures using the GAB and BET sorption models (Figs. 2 and 3). Also, linear relationships were established for the Oswin and Smith sorption models (Figs. 4 and 5). The magnitude of the  $a_w/M_e$  increases with the increase in the storage temperature until 0.6  $a_w$  when it starts decreasing for the GAB models. The BET models, however, show a continuous s increase in the magnitude of the  $a_w/(1-a_w)M_e$  with an increase in the storage temperature and  $a_w$ . An increase in the equilibrium moisture content with an increase in the storage temperature and  $a_w$  was observed for the Oswin and Smith models. The observed differences in the variable interaction may be due to the manner in which the models were constituted for sorption isotherm description. A similar result was reported by Arevalo et al. (2004) in their work on mathematical modelling of Inga pulp. Also, the research work of Oyelade et al. (2008) on the sorption isotherm of maize flour corroborates the findings of this study.




**Fig 1.** Adsorption Isotherm Curve of *Pupuru* flour at different temperatures




**Fig 2.** Sorption isotherm of stored *Pupuru* flour based on GAB model



**Fig 3.** Sorption isotherm of stored *Pupuru* flour based on BET model



**Fig 4.** Sorption isotherm of stored *Pupuru* flour based on Oswin model



**Fig 5.** Sorption isotherm of stored *Pupuru* flour based on Smith model

The parameters of the sorption modelling of the *pupuru* flour at different storage temperatures are shown in Table 2. The magnitudes of the  $\alpha$ ,  $\beta$  and  $\gamma$  represent the coefficients of the terms of the models used for the sorption isotherm prediction. The values of the constants  $\alpha$ ,  $\beta$  and  $\gamma$  obtained from the curves (Abramović and Klofutar, 2006) were used to calculate the three model parameters of  $C$ ,  $K$  and  $M_o$ . The monolayer ( $M_o$ ) values obtained were temperature dependent, and this agrees with previous studies (Abramović and Klofutar, 2006; Fadeyibi et al., 2012). The  $M_o$  was found to decrease with an increase in the temperature for GAB, BET, and Oswin models. It is important to note that an increase in the temperature usually leads to a decrease in  $M_o$  (Arevalo-pinedo et al., 2004) and this was observed for the GAB, BET and Oswin models. Thus, the decrease in the  $M_o$  may indicate that the adsorbed molecules will require less kinetic energy to break away from their sorption sites. The values obtained were consistent with other similar unenriched cassava products such as *fufu* (0.043–0.049), tapioca (0.049–0.058) and unenriched *gari* (0.057–0.060) (Kuye and Sanni, 2002; Oluwamukomi et al., 2008). The  $M_o$  was less than 0.1 kg kg<sup>-1</sup> (db) in all the

samples, which was the maximum value reported for the most food materials (Tejada-Ortigoza et al., 2017; Oluwamukomi et al., 2008; Kaymak- Ertekin and Sultanoglu, 2001). These values agree with literature values for other starchy products (Kuye and Sanni, 2002; Oluwamukomi et al., 2008). On the contrary, the  $M_o$  estimated from the Smith model increases with an increase in the temperature, and this may require high energy to process the product. Furthermore, the parameter  $C$  was found to increase with an increase in the storage temperature for the GAB model, but decrease for the BET model. However, the parameter  $K$  decreases with an increase in the temperature for the GAB and the Smith models, but remains inconsistent for the Oswin model. The decrease in the  $K$  values may indicate that the multilayer molecules of the *pupuru* flour may require less entropy for processing at higher temperature. This will usually result in the molecules requiring less kinetic energy to cause them to pull apart (Abramović and Klofutar, 2006; Oluwamukomi et al., 2008). Thus, the GAB, BET and Oswin models are better estimators of the  $M_o$ , while the parameters  $C$  and  $K$  were better estimated using the GAB model.

**Table 2.** Parameters of sorption modelling of *Pupuru* flour at different storage temperatures

Sorption Isotherm Model																	
GAB				BET				Oswin				Smith					
Temperature (°C)			MP	Temperature (°C)			MP	Temperature (°C)			MP	Temperature (°C)			MP		
α	β	γ		α	β	γ		α	β	γ		α	β	γ			
-14.3	17.5	0.41	10	-13.5	16.7	0.10	50	63.26	59.69	41.85	-2.36	-2.256	-1.942	-0.067	-0.067	-0.098	
-9.93	12.29	0.044	30	-13.83	-13.3	0.000	50	-9.13	0.313	0.286	0.299	0.046	0.054	0.069			
0.06	0.05	0.041	10	0.000	0.000	0.000	30	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	
55.3	219.4	348.7	50	-3.574	-3.68	-5.54	10	0.020	0.313	0.286	0.279	0.046	0.054	0.069			
-19.0	-23.3	-26.4	0.000	0.00	0.000	0.000	0.000	-2.36	-2.256	-1.942	-0.067	-0.067	-0.098				

*k*, *c*, are model constants; MP is model parameter;  $M_o$  is monolayer moisture content;  $\alpha$ ,  $\beta$  and  $\gamma$  are regression coefficients

### Validation of Sorption Data


The results of the model verification and calibration are shown in Table 3. The sorption data adequately fit the GAB and Smith models followed by the Oswin model and the least for the BET model. This is evident from the differences in the values of the  $R^2$  and  $R_{adj}^2$  of the regression equations obtained from the analysis of the 4 models. This means that the GAB, BET and Smith models can be used for describing the sorption behaviour of the *Pupuru* flour over the entire range of the  $a_w$ . The Oswin model may not be good for predicting the sorption isotherm of the *pupuru* flour because of its lower  $R^2$  and  $R_{adj}^2$  values relative to the

other models tested. Also, the relationship between the actual and model predicted equilibrium moisture contents of the product is shown in Fig. 6. The predicted values closely match the actual values of the equilibrium moisture content of the product. All the models are adequate and valid for independent prediction of the equilibrium moisture content since the MSE values are generally less than 10% at the storage temperatures considered. This is consistent with the findings of Oluwamukomi et al. (2008) for unenriched *gari* product and Kuye and Sanni (2002) for *lafun* food product. Thus, the model can be used for predicting the sorption isotherm behaviour of the *pupuru* flour within its range of  $a_w$ .

**Table 3.** Verification data of the sorption models for *pupuru* flour at different temperatures

Sorption Isotherm Model												
GAB			BET			Oswin			Smith			
Temperature (°C)			Temperature (°C)			Temperature (°C)			Temperature (°C)			
10	30	50	10	30	50	10	30	50	10	30	50	
$R^2$	0.985	0.98	0.977	0.738	0.743	0.774	0.980	0.959	0.981	0.997	0.996	0.994
$R_{adj}^2$	0.981	0.975	0.971	0.673	0.679	0.718	0.975	0.949	0.976	0.996	0.995	0.993
MSE	0.011	0.007	0.002	0.006	0.005	0.005	0.005	0.004	0.001	0.004	0.003	0.006

MSE is mean square error,  $R^2$  is coefficient of determination and  $R_{adj}^2$  adjusted coefficient of determination



**Fig 6.** Relationship between the actual and model predicted equilibrium moisture content of stored *pupuru* flour based on (a) GAB model (b) BET model (c) Oswin model (d) Smith model

## Conclusion

This study was carried out to describe the sorption isotherm behaviour of the *pupuru* flour at different temperatures and water activity conditions. The sorption data were fitted into the GAB, BET, Oswin and Smith models for starchy foods. The model parameters were found to vary with an increase in the storage temperature and the  $a_w$  of the product. The sorption parameters ( $M_o$ ,  $C$  and  $K$ ) were generally higher for the products stored at 10 °C and 0.10  $a_w$ . The  $M_o$  was found to decrease with an increase in the temperature for GAB, BET, and Oswin models. On the contrary, the  $M_o$  estimated from the Smith model increases with an increase in the temperature, and this may mean high energy is required to process the product. The parameter  $C$  was found to increase with an increase in the storage temperature for the GAB model, but decrease for the BET model. However, the parameter  $K$  decreases with an increase in the temperature for the GAB and the Smith models, but remains inconsistent for the Oswin model. The high values of the  $R^2$  and  $R^2_{adj}$ , and the  $MSE < 10\%$  indicate the ability of the models to predict the sorption isotherm characteristics of the *pupuru* flour within the range of  $a_w$  and the storage temperatures considered. The GAB, BET and Oswin models are better estimators of the  $M_o$ , while the parameters  $C$  and  $K$  were better estimated using the GAB model.

**Author Contributions:** Dr. Adeshina Fadeyibi conceived the research idea, analysed the data, interpreted the results and drafted the final manuscript. Dr. Janet Oluwatoyin Alaba conducted the research, analysed the data and drafted the original draft.

**Funding:** This research received no external funding.

**Acknowledgments:** The authors acknowledge the technical assistance provided by the staff of the Food Science and Laboratory of the Wesley University of Science and Technology, Ondo State, Nigeria.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

Abramović, H., Klofutar, C. (2006): Water adsorption isotherms of some gellan gum samples. *Journal of food engineering* 77 (3), 514-20.

Adebowale, A.A., Owo, H.O., Sobukola, Obadina, O.A., Sanni, L.O., Awonorin, S. O. (2017): Influence of Storage Conditions and Packaging Materials on Some Quality Attributes of Water Yam Flour Cogent. *Food and Agricultural* 3 (1), 1-26.

Aguirre-Loredo, R.Y., Rodriguez-Hernandez, A.I., Velazquez, G. (2017): Modelling the effect of temperature on the water sorption isotherms of chitosan films. *Food Science and Technology* 37 (1), 112-118.

Ajisegiri, E.S.A., Chukwu, O., Sopade, P.A. (2007): Moisture- Sorption Study of Locally Parboiled Rice. *Australian Journal of Technology* 11 (2), 86-90.

AOAC (2005): Official Method of analysis. (Helrich, ed.) 15th edition., Association of Official Analytical Chemists, Arlington, VA. Vol. 1.

Arevalo-Pinedo, A. D., Giraldo-Zuniga, F. L., Santos, Z. D., Arevalo, S., Rosalinda, P. A. (2004): Application of Mathematical models of two and three parameters in the prediction of sorption isotherms for Inga (Inga edulis) pulp. Drying 2004-Proceedings of the 14th International Drying Symposium (IDS 2004) Sao Paulo, Brazil, 22-25, 628-633.

Bello, K., Sarojini, B.K., Narayana, B., Rao, A., Byrappa, K. (2018): A study on adsorption behavior of newly synthesized banana pseudo-stem derived superabsorbent hydrogels for cationic and anionic dye removal from effluents. *Carbohydrate polymers* 181, 605-15.

Cova, A., Sandoval, A.J., Balsamo, V., Müller, A.J. (2010): The effect of hydrophobic modifications on the adsorption isotherms of cassava starch. *Carbohydrate Polymers* 81 (3), 660-667.

Daramola O., Idowu, M., Atanda, O., Oguntona, C. (2010): Effects of Packaging material on the quality of "pupuru" flour during storage. *African Journal of Food Science* (4), 258-263.

Erbaş, M., Candal, C., Kılıç, Ö., Mutlu, C. (2016): Determination and solution of moisture sorption isotherms of foods." *GIDA-Journal of Food* 41 (3), 171-178.

Fadeyibi, A., Osunde, Z. D., Ussaini, M. S., Idah, P. A., Balami, A. A. (2012): Evaluating monolayer moisture content of rubber seed using BET and GAB sorption equations. *International Journal of Farming Allied Sciences* 1, 72-76.

Famurewa, J.A.V., Oluwamukomi, M.O., Alaba, J.O. (2012): Storage stability of *pupuru* flour (a cassava product) at room temperature. *British Journal of Applied Science and Technology* 2 (2),138.

Ilouno, B.E.L., Ndimele, E.C., Adikwu, O., Obiekezie, S.O. (2016): Bacteriological quality of some fermented food products in Keffi, Nasarawa state. *FULafia Journal of Science and Technology* 2 (1), 4-10.

Kaymak-Ertekin F., Sultanoglu, M. (2001): Moisture sorption isotherm characteristics of peppers. *Journal of Food Engineering* 47, 225-231.

Kumar A.J., Singh R.R., Patil G.R., Patel A.A. (2005): Effect of temperature on moisture desorption isotherms of kheer. *LWT-Food Science and Technology* 38 (3), 303-310.

Kuye, A., Sanni, L.O. (2002): Analysis of the equilibrium moisture sorption data for lafun and soyflour. *Journal of Modeling, Design and Management of Engineering Systems* 1 (1), 63-70.

Labuza, T.P. and Altunakar, B. (2020): Water activity prediction and moisture sorption isotherms. *Water activity in foods: fundamentals and applications* 61-205.

Mitrevska, V., Mitrevska, C., Mijakovski, V., Pavkov, I.S., Geramitcioski, T. (2017): Mathematical modelling of the sorption isotherms of quince. *Thermal Science* 21 (5), 1965-1973.

Ojuko O.O., Olapade A.A. (2018): Effect of Addition of Yellow-Fleshed Sweet Potato Flour (*Ipomoea batatas*) and Fermentation on Chemical, Functional, Microbial and Sensory Attributes of an Instant Fura Powder. *Journal of Scientific Research and Reports* 10 (1), 1-2.

Oluwamukomi, M.O., Adeyemi, I.A., Odeyemi, O.O. (2008): Adsorption Isotherm and Thermodynamic Characteristics of Soy-Melon Enriched "Gari" Semolina from Cassava". *Agricultural Engineering International: CIGR E journal*. FP 07010(10).

Osundahunsi, O.F. (2005): Effect of drying methods on composition, sensory evaluation and rheological value of *pupuru* (fermented cassava product). *Journal of Food Technology* 3 (3), 353-355.

Oyelade, O.J., Tunde-Akintunde, T.Y., Igbeka, J.C., Oke, M.O., Raji, O.Y. (2008): Modelling moisture sorption isotherms for maize flour. *Journal of Stored Products Research* 44 (2), 179-185.

Rockland, L.B., Stewart, G.F. (2013): Water activity: influences on food quality: a treatise on the influence of bound and free water on the quality and stability of foods and other natural products. Academic Press; 2013 Oct 22.

Samapundo, S., Devlieghere, F., De Meulenaer, B., Atukwase, A., Lamboni, Y., Debevere, J.M. (2007): Sorption isotherms and isosteric heats of sorption of whole yellow dent corn. *Journal of Food Engineering* 79 (1), 168-175.

Shittu, T. A, Lasekan, O.O. Sanni, L.O., Oladosu, M.O. (2001): The effect of drying methods on the functional and sensory characteristics of *pukuru*- a fermented cassava product. *ASSET Series A* 1 (2), 9-16.

Shittu, T.A., Oyewole, O.B., Olawuyi, O., Daramola, O. (2003): Processing Technology of *Pupuru*: A survey of Practices and Product Quality in the South West of Nigeria. *ASSET Series B* 2 (2), 17-27.

Singh, P., Talukdar, P. (2020): Determination of Desorption Isotherms of Potato Using Gravimetric Method and Fast Isotherm Method. *Heat Transfer Engineering* 41 (6-7), 513-521.

Tejada-Ortigoza, V., Garcia-Amezquita, L.E., Serment-Moreno, V., Torres, J.A., Welti-Chanes, J. (2017): Moisture sorption isotherms of high pressure treated fruit peels used as dietary fiber sources. *Innovative Food Science and Emerging Technologies*, 45-53.

Yan, Z., Sousa-Gallagher, M.J., Oliveira, F.A. (2008): Sorption isotherms and moisture sorption hysteresis of intermediate moisture content banana. *Journal of Food Engineering* 86 (3), 342-8