



Original scientific paper

DOI: 10.17508/CJFST.2021.13.2.10

Antioxidant, Nutritional, and Physicochemical Quality of Yoghurt Produced from a Milk-Based Fermentation Mix Enhanced with Food Spices

 Oludare Ogunyemi^{1,2*}, Gideon Gyebi¹, Raheem Shaibu¹, Mercy Fabusiwa¹, Charles Olaiya²

¹Human Nutraceuticals and Bioinformatics Unit, Department of Biochemistry, College of Natural and Applied Sciences, Salem University, Lokoja, Nigeria

²Nutritional and Industrial Biochemistry Research Unit, Department of Biochemistry, College of Medicine, University of Ibadan, Ibadan 200005, Nigeria

ARTICLE INFO

Article history:

Received: July 7, 2020

Accepted: March 23, 2021

Keywords:

spices
functional foods
antioxidants
sensory quality
yoghurt

ABSTRACT

Utilization of natural additives, which could meet the growing need for 'clean label' and value addition with respect to disease prevention and health promotion in humans, is a current interest in the yoghurt industry. In this study, yoghurt recipes derived from a standard milk-based fermentation mix enhanced with African black pepper (ABP), turmeric, or cloves extracts were evaluated for antioxidant, nutritional and physicochemical qualities. Each fermentation mix was pasteurized, inoculated with 0.5% industrial starter culture containing *Streptococcus thermophilus* and *Lactobacillus delbrueckii* subsp. *bulgaricus* (1:1), and then allowed to ferment in a fermentation tank for 9 hours at 40 °C. The results revealed that the clove-enriched yoghurt recipe exhibits the strongest antioxidant capacity, as indicated by a significant increase ($p < 0.05$) in total phenolics, total flavonoids, total antioxidant capacity (TAC), DPPH radical scavenging activity, nitric oxide radical scavenging activity, ferric ion reducing power (FRAP), and inhibition of lipid peroxidation, but it also exhibits reduced apparent viscosity and organoleptic properties. Similarly, ABP increased the antioxidant characteristics, with increased protein content and gel viscosity. Turmeric extracts significantly increased ($p < 0.05$) the calcium content and reduced syneresis. In conclusion, the fermentation of a milk-based mix containing ABP extract resulted in yoghurt which combines improved antioxidant characteristics with increased protein and enhanced techno-functionality.

Introduction

Yoghurt is globally regarded as a functional food which combines unique nutritional values with the promotion of gut health, heart functions, and the natural immune defence in humans (Hashemi Gahruei et al., 2015; Caleja et al., 2016; Tavakoli et al., 2019). During the fermentation of milk, lactic acid bacteria (LAB) utilize lactose and nitrogenous compounds leading to the production of extracellular lactic acid and various other primary metabolites that provoke acidification of the medium. This causes several chemical changes, which include a decrease of pH, increased nutrient

bioavailability, formation of flavour compounds, coagulation of proteins, and subsequent gel formation (Tamang et al., 2016; Brückner-Gühmann et al., 2019). In the acidic medium, the net negative charge on the casein micelles decreases, and thereby reduces electrostatic repulsion between charged groups, causing coagulation and destabilization of the casein micelles and the conversion of the fluid milk into a viscoelastic gel (Brückner-Gühmann et al., 2019). These changes greatly influence the physicochemical characteristics and consumers' acceptability of yoghurt. While the texture components of yoghurt can be maintained and improved by optimizing the production process, the flavour, shelf-

*Corresponding author E-mail: omogunyemi1@gmail.com

life, and health benefits can be modified or enhanced by the utilization of additives (Jaworska et al., 2005; Brückner-Gühmann et al., 2019).

A current interest in the yoghurt industry is the utilization of natural additives towards developing new products which do not only cater to consumers' expectations regarding nutritional value and techno-functionality, but also meet the growing need for 'clean label' and value addition with respect to antioxidant potentials, disease prevention, and health promotion in humans. In this direction, herbal yoghurts, which are enhanced with *Lycium barbarum* extract (Baba et al., 2014), tea extracts (Muniandy et al., 2016), and mangosteen rind extract (Wibawanti et al., 2018) have been reported earlier. Antioxidant components of foods play a significant role in disease prevention and the maintenance of health and wellbeing. They help in the deactivation of free radicals which can cause cell and tissue damage, leading to various diseases of the modern human. Epidemiological studies have shown that antioxidants can prevent the development of degenerative diseases such as cancer, coronary heart diseases, obesity, type 2 diabetes, hypertension, premature aging, and inflammatory diseases (Senadeera et al., 2018). Aside from that, antioxidant compounds may enhance protection against lipid peroxidation and peroxidability of yoghurt, thereby improving the shelf-life of the fermented product. Antioxidant capacity assays of food substances are useful in measuring the overall antioxidant activity due to the contents of the antioxidant compounds in the food matrix. Flavonoids constitute a large group of polyphenolic phytochemicals with antioxidant properties which are overwhelmingly exerted through direct free radical scavenging as reducing agents, hydrogen donors, and singlet oxygen quenchers (Beckman, 2000). Flavonoids also exhibit antioxidant properties through chelating with transition metals, primarily Fe(II), Fe(III), and Cu(II), which participate in reactions generating free radicals in a biological system. Interestingly, the metal-flavonoid chelates formed are even more potent free radical scavengers than the parent flavonoids and play a prominent role in protection against oxidative stress (Dusan and Vesna, 2007; Cherrak et al., 2016). These antioxidant compounds are of interest in the food industry and human nutrition because of their attractive colour, stability in high acid foods, and antioxidant power (Panche et al., 2016).

Food spices have multifarious roles such as antioxidant, medicinal, food flavouring, and preservative functions (Olaiya et al., 2013; Kapadiya et al., 2016; Gyebi et al., 2019). Our preliminary assessment revealed that the addition of extracts from several spices to a milk-based fermentation mix improved the antioxidant potentials of yoghurt in a concentration-dependent manner and also increased the consumer acceptability of the products

(Ogunyemi et al., 2020). The current study focused on further evaluation of the antioxidant, nutritional, and physicochemical characteristics of yoghurt enriched with 4% extracts of African black pepper (ABP) (*Piper guineense*), turmeric (*Curcuma longa*), and cloves (*Syzygium aromaticum* L.).

Materials and methods

Preparation of spice extracts

The spice extracts were prepared from African black pepper (ABP), turmeric, and cloves as reported previously (Ogunyemi et al., 2020). The spices were purchased from the International Market, Lokoja, Kogi State, Nigeria. They were washed several times with tap water (portable water), then peeled, rewashed with tap water, sliced into small sizes of 2-3 diameter thickness and dried in a vacuum oven at 60 – 65 °C for 72 hours. The dried spices were blended into powder using a kitchen blender. The powder obtained was extracted with water to obtain the spice extracts.

Preparation of spiced yoghurt samples

Commercial skimmed milk comprising of 32% protein, 54% lactose, 0.5% milk fat, and 9% minerals was procured from DANO milk, Nigeria. An industrial freeze-dried starter culture (YoFlex CHR Hansen Company, Denmark) containing *Streptococcus thermophilus* and *Lactobacillus delbrueckii* subsp. *bulgaricus* (1:1) was purchased. The samples of yoghurt were produced according to the International Standard of yoghurt as described by Guler and Mutlu (2005). Here, 400 g of skimmed powdered milk (comprising of 32% protein, 54% lactose, 0.5% milk fat, and 9% minerals) was reconstituted with water and heated to 80 °C for 15 minutes for pasteurization, and then allowed to cool to 42-45 °C before inoculation with the starter culture. The milk mixture was divided into four portions, plain yoghurt, ABP-flavoured, turmeric flavoured, and clove spiced. They were incubated at 40 °C and the fermentation process was monitored for 9 hours via the production of lactic acid and the pH value, until a pH of about 4.9 was attained, as reported in previous studies (Ogunyemi et al., 2020). The yoghurt samples were then packed and stored at 4 °C until further analysis.

Antioxidant analysis

Total phenolic content in yoghurt samples was determined using the Folin Ciocalteu reagent (FCR) method described previously by Singleton et al. (1999). The flavonoids were measured by aluminium chloride colorimetric assay as described by Zhishen et al. (1999).

The DPPH• assay was carried out as described by Shirwaikar et al. (2006). Nitric oxide radical scavenging activity was determined according to the method reported by Johnson (1964). The ability of the samples to reduce oxidized iron was carried out as defined by Oyaizu (1986). The total antioxidant capacity (TAC) assay was carried out as described by Re et al. (1999). The inhibition of lipid peroxidation was carried out in accordance with the methods described by Ruberto et al. (2000) with slight modification.

Chemical and physicochemical analysis

Protein, ash, moisture content, and total solids (TS) of the yoghurts were determined according to Latimer and the Association of Official Analytical Chemists (2019). The pH-value and total titratable acidity (TTA) were measured during fermentation. Titratable acidity in terms of lactic acid was measured according to Tomovska et al. (2016). The amount of NaOH used up was recorded and TTA was calculated as shown below.

$$\text{Titratable acidity} = (\text{Titre value} \times 0.009) / (\text{Weight of sample}) \times 100 \quad (1)$$

The apparent viscosity of the homogenized samples was measured using a Brookfield viscometer (Brookfield Programmable Rheometer, Model RVDV-III Ultra; Brookfield Engineering Laboratories, Stoughton, MA, USA) with a spindle no. 4 and the rotation speed of 10 rpm at 24 °C ± 1 °C. The values of apparent viscosity were recorded in centipoises (cP). Syneresis of the different yogurt samples was determined according to the methodology proposed by Dabija et al. (2018). 100 mL of each sample was placed in a funnel lined with Whatman filter paper number 1. After 6 h of drainage, the volume of whey was measured and the susceptibility of syneresis was calculated thus:

$$\text{Syneresis} = (\text{volume of whey collected after drainage}) / (\text{volume of yogurt sample}) \times 100 \quad (2)$$

The colour of yoghurt samples was determined by the Minolta colour meter CR-410 model (Minolta Co., Osaka, Japan). The calibration of the meter was achieved using a white standard calibration plate ($L^* = 92.95$, $a^* = -4.86$, $b^* = -6.65$). The CIE L^* , a^* , b^* and ΔE values of the yoghurt were obtained directly from the meter. The hue angle (h^*) and chroma (C^*) were calculated.

Sensory evaluation

A total of 20 panellists from the Salem University Nigeria community, including university students and staff, participated in the study. The panellists were

chosen based on their willingness, availability, and motivation. Sensory evaluation of plain and spice yoghurt was carried out to determine their organoleptic characteristics in terms of their colour, aroma, taste, texture, and overall acceptance. A nine-point hedonic scale, varying from dislike extremely (score 1) to like extremely (score 9) was used (Ogunyemi et al., 2020). To avoid bias, samples were coded so that the panellists could not identify them. The sample presentation order was randomized among and within the assessors.

Statistical analysis

The data generated from the study was subjected to analyses using the SAS 9.12 version and Graph Pad Prism. One-way analysis of variance (ANOVA) was used to detect the treatment effects. The means were separated by the Duncan multiple range test (DMRT) and a probability value less than 0.05 was considered statistically significant.

Results

Fermentation of spiced yoghurt

The fermentation process of the spiced yoghurt samples was monitored using the decrease in pH and lactic acid accumulation as reported in figure 1.




Fig. 1. Effects of spice extracts on pH reduction (a), and total acid accumulation (b), of milk during fermentation.

Addition of the spice extracts had only slight effects on the fermentation process with respect to decreasing the pH (Figure 1a), as compared to that of the plain yoghurt (control). Similar results were recorded with respect to lactic accumulation (Figure 1b). The pH of the final yoghurt samples ranged between 4.9 and 5.1.

Antioxidant potentials of the spiced yoghurts

Figure 2 shows the in vitro antioxidant potentials of spiced yoghurt samples.




Fig. 2. Antioxidant potentials of spiced yoghurt; The means of rank of classification of yoghurt samples are represented in alphabets; values with the same letter in the same column are not significantly different ($P>0.05$), (a) total phenolic and total flavonoid contents (b) ferric reducing antioxidant capacity (FRAP), total antioxidant capacity (TAC), (c) DPPH radical scavenging activity, nitric oxide scavenging activity and lipid peroxidation. ABP = African black pepper.

Even though a small amount of phenolics (9.78 ± 0.57 mg GAE/100mL) was detected in the plain yoghurt, this amount significantly increased ($p<0.05$) in all the spiced yoghurt samples. The addition of clove extracts resulted in a 0.72-fold increase in phenolic content as compared to the plain yoghurt (Figure 2a). Similar effects were also recorded for total flavonoid content in the yoghurt samples (Figure 2a). Cloves gave the highest level of TAC and FRAP, as observed in Figure 2b and 2c. The results also revealed that most spice extracts significantly increased ($p<0.05$) the inhibitory activity of yoghurt samples against the DPPH radical. The addition of 4% clove extract gave the highest inhibitory activity, with up to 0.62-fold increases in DPPH scavenging activity as compared with the plain yoghurt. The nitric oxide radical scavenging activity of the yoghurt samples as shown in Figure 2c also shows a similar pattern with the DPPH scavenging assay. Both the plain yoghurt and spiced yoghurt also exhibit inhibition of lipid peroxidation and most of the spiced yoghurt had significantly higher inhibition than the plain yoghurt (figure 2c).

Effects of spices on some nutrients in spiced yoghurt

Table 1 shows the effects of the incorporation of spice extracts on some nutrients of the yoghurt. While supplementation of yoghurt with the spice extracts had little or no effects on the moisture contents, all spice extracts significantly increased ($p < 0.05$) the protein content of the yoghurt samples as compared to the plain yoghurt. Clove extracts gave the highest increase in protein content with a 0.61-fold significant increase, followed by turmeric and then ABP. Regarding ash content, while the addition of ABP and turmeric had little or no effect on the yoghurt samples, the clove extract significantly increased the ash content. While the addition of clove extracts had little or no effect on the calcium content, turmeric significantly increased ($p < 0.05$) the content of calcium in the yoghurt sample as compared with the plain yoghurt.

Physicochemical characteristics of spiced yoghurt samples


The results shown in figure 4 revealed the effects of spice addition on the viscosity of the spiced yoghurt samples.

The addition of ABP increased the apparent viscosity as compared to the plain yoghurt and the other spiced yoghurt samples. The addition of turmeric and cloves reduced the values of apparent viscosity.

Table 1. Effects of 4% spice extracts on the contents of some nutrients in yoghurt

Yoghurt	Moisture (%)	Protein (%)	Ash (%)	Ca (mg/100 mL)	P (mg/100 mL)
Control	79.46 ± 1.70 ^a	2.54 ± 0.12 ^b	11.28 ± 0.53 ^{ab}	610.67 ± 5.21 ^{ab}	2159.83 ± 81.67 ^a
ABP	82.09 ± 0.88 ^a	3.67 ± 0.19 ^a	9.32 ± 0.56 ^b	462.67 ± 93.93 ^b	1549.67 ± 57.76 ^b
Turmeric	82.38 ± 0.19 ^a	3.80 ± 0.23 ^a	10.34 ± 0.36 ^{ab}	682.50 ± 58.97 ^a	1215.10 ± 98.84 ^c
Clove	82.58 ± 0.30 ^a	3.96 ± 0.16 ^a	12.40 ± 1.50 ^a	635.83 ± 32.73 ^{ab}	1120.77 ± 15.12 ^c

Mean values (N=3) with different superscript in each column are significantly (p<0.05) different from one another. The means were separated by the Duncan multiple range test (DMRT) and a probability value less than 0.05 was considered statistically significant ABP = African black pepper

**Fig. 3.** Apparent viscosity of yoghurt samples measured in centipoises (cP).**Table 2.** Effects of spice extracts on physicochemical properties of yoghurt

Parameters	Control	ABP	Turmeric	Clove
pH	4.82 ± 0.05 ^a	4.79 ± 0.02 ^a	4.76 ± 0.01 ^a	4.66 ± 0.12 ^a
Total Acid (%)	7.82 ± 0.05 ^a	7.65 ± 0.09 ^a	6.48 ± 0.16 ^b	7.65 ± 0.18 ^a
TSS (Brix)	14.75 ± 0.58 ^a	13.50 ± 1.01 ^{ab}	12.83 ± 0.36 ^{ab}	12.17 ± 0.58 ^b
Syneresis	64.00 ± 1.15 ^a	65.33 ± 0.67 ^a	53.33 ± 3.53 ^b	59.33 ± 1.76 ^b
WHC	80.00 ± 1.15 ^a	74.00 ± 1.53 ^a	76.33 ± 3.18 ^a	73.33 ± 1.76 ^a
L*	70.54 ± 0.11 ^a	66.62 ± 2.58 ^a	66.90 ± 4.71 ^a	66.61 ± 6.24 ^a
a*	-13.37 ± 0.09 ^a	-12.40 ± 0.59 ^a	-13.44 ± 0.99 ^a	-13.00 ± 2.23 ^a
b*	11.25 ± 0.16 ^a	11.18 ± 0.56 ^a	13.61 ± 1.22 ^a	12.76 ± 1.14 ^a
H*	-0.69 ± 0.00 ^a	-0.73 ± 0.01 ^a	0.79 ± 0.03 ^a	0.79 ± 0.05 ^a
C*	17.46 ± 0.18 ^a	16.69 ± 0.80 ^a	19.15 ± 1.48 ^a	18.27 ± 2.31 ^a
ΔE	45.33 ± 1.76 ^a	41.03 ± 2.51 ^a	31.05 ± 9.64 ^a	38.71 ± 6.97 ^a


Mean values (N=3) with different superscript in each column are significantly (p<0.05) different from one another. The means were separated by the Duncan multiple range test (DMRT) and a probability value less than 0.05 was considered statistically significant

The addition of both turmeric and clove extracts significantly reduced the syneresis of the yoghurt samples by 0.45-fold and 0.48-fold, respectively. The addition of spice extracts to yoghurt did not have a significant effect on the colour characteristics of the yoghurt (Table 2). The results in table 2 revealed that the enrichment of the milk-based fermentation mix with the spice extracts did not cause significant change to the colour characteristics of the yoghurt samples in terms of the degree of lightness (L*), red-green range (a*), yellow-blue range (b*), chroma (C*), hue angle (h), and the total colour difference (ΔE), compared to the control.

Sensory Characteristics

Consumer acceptability is a key factor in product development. The results of the sensory profile

analysis of the spiced yoghurts are summarized in Figure 4.

**Fig. 4.** Sensory characteristics of spiced yoghurt samples. The means of rank of classification of yoghurt samples are

represented in alphabets; values with the same letter in the same column are not significantly different ($p>0.05$).

Discussion

The enrichment of a milk-based fermentation mix with fruit, food spices, and medicinal herbs as additives presents a novel approach for developing functional yoghurts with improved nutritional, antioxidant, and therapeutic properties. The addition of the spice extracts used in this study had little or no effect on the fermentation process as monitored with the decrease in pH and the accumulation of lactic acid. The fermentation of the plain and spiced yoghurt may be said to follow the general fermentation process (Figure 1) which occurs in three phases: the lag phase characterized by a slow pH decrease, the logarithmic phase with a rapid pH decrease and eventual decelerated acidification (Corrieu and Béal, 2016). This is generally accompanied by lactic acid accumulation and the coagulation of milk beginning at pH 5.4 and completing at pH 4.6 (Güner et al., 2007; Lee and Lucey, 2010).

The endogenous antioxidant potential exhibited by the plain yoghurt in this study may be due to substances such as peptides and free amino and fatty acids with antioxidant activities that are generated from the milk during fermentation as reported earlier by Tavakoli et al. (2019). A biochemical event such as proteolysis of milk proteins may yield amino acids with phenolic side chains such as tyrosine which react with the Folin-Ciocalteu reagent (Raikos et al., 2018). Also, the metabolism of phenolic compounds by the lactic acid bacteria may be associated with flavonoid glycoside hydrolysis or the C-ring cleavage and the release of simple phenolics such as phenolic acids (Muniandy et al., 2016). The enhancement of the antioxidant potentials of yoghurt samples with the spice extracts may be due to the enrichment of the yoghurt with phenolic and flavonoid compounds, as these spices are known as good sources of polyphenolic compounds (Kapadiya et al., 2016). Furthermore, some endogenous peptides may also be able to exert a marked synergistic effects with phenolic antioxidants from plant products (Citta et al., 2017). The influence of the spice extract on the antioxidant capacity of the yoghurt samples is consistent with earlier reports on supplementation of cow, buffalo, and goat milk yogurts with aqueous extracts of *Zingiber officinale* and *Beta vulgaris* (Srivastava et al., 2015). These authors reported that the highest antioxidant activities with DPPH and FRAP methods were found in 2% ginger rhizome goat milk yoghurt and 2% beet root goat milk yoghurt, followed by 2% ginger cow milk yoghurt. The lowest antioxidant activity was found in

buffalo milk herbal yoghurt. Another study incorporated vegetable extracts (carrots, pumpkin, broccoli, and red sweet pepper) at 10% concentration and ferric reducing antioxidant power (FRAP), and the DPPH assays were used for anti-oxidant activity during the storage period of 14 days. Yoghurt with broccoli and red sweet pepper revealed higher DPPH free radical scavenging activity and FRAP (Najgebauer-Lejko and Sady, 2015). Yoghurt was supplemented with fruit pulp of papaya and cactus pear using *Lactobacillus bulgaricus* and *Streptococcus thermophiles* as starter cultures, and total phenolic contents, ascorbic acid, and total antioxidant activity were analysed. Yoghurt with the addition of papaya fruit pulp had higher total phenolic contents, antioxidant activity, and Vitamin C concentration (Khan et al., 2019).

The moisture content of the spiced yoghurt was maintained within the range of most yoghurt available in the market (80-86%) (Ndife, 2014). Enrichment of the yoghurt samples with plant proteins from the spices might account for the improved protein content of the spiced yoghurt. It is known that the addition of plant proteins leads to the improvement of the quality of yogurt and an increase in its nutritional value (Dabija et al., 2018). Many researchers have successfully fortified yogurt with plant proteins (Akalin et al., 2012; Morell et al., 2015). Such fortification may have positive impacts on the textural and physical properties of yogurts such as yogurt firmness, viscosity, and functional properties. The firming effect of the addition of different proteins in the formulation of yogurts could lead to a distinctive structure of casein, with desirable effects on the textural characteristics of the finished product (Dabija et al., 2018). On the other hand, for the consumer, the increase in the protein content improves the degree of satiety, especially beneficial for those who consume fermented dairy products for health reasons, certain diets, etc. (Dabija et al., 2018). Ash content is a reflection of the amount of minerals in a food material (Gemedé et al., 2015). Calcium, which plays key roles in bone formation and mineralization, is highly required during growth, pregnancy, and lactation (Soetan et al., 2009; Gemedé et al., 2015). This, therefore, means that turmeric enriched yoghurt would be a beneficial drink for children, pregnant and lactating mothers, as well as the elderly, whose calcium requirement is high.

Supplementation of yoghurt with ABP gave the highest viscosity as compared to the plain yoghurt and other spiced yoghurt samples. This is in contrast with the general belief that the addition of plant extracts decreases the consistency of the products, owing to reduced water-binding capacity of proteins

(Ramaswamy and Basak, 1992; El-Said et al., 2014). The improved viscosity may be due to the contribution of protein (as reported in table 4) to the fermentation medium, leading to the improvement of the technofunctionality of the product. This can be attributed to the positive impact of protein enrichment on the aggregation of the casein network in yoghurts via electrostatic interaction, and on the resistance of the flow of the yoghurt matrix. This study also shows that the addition of turmeric and clove extracts helped in the reduction of syneresis of the spiced yoghurt. Tendency to syneresis gives information about the stability of the gel samples. The rate of syneresis in acidified milk gels like yoghurt is mainly determined by the microstructure of the protein network. If the water binding is not sufficient, whey will be expelled on the surface of the product during storage. During the yogurt making process, the level of lactic acid becomes higher as fermentation goes on, and a series of cascades such as an increase in acidity, dissociation of the carboxyl groups, ionization of serine phosphate, and the increase in the negative charge between casein micelles take place, respectively. At this point, whey proteins are denatured and are more susceptible to association with casein and casein micelles. Additionally, during acidification, the denatured whey proteins, associated or not associated with casein micelles, aggregate. This is the result of the neutralization between calcium phosphate and the increasing negative charge, and due to the presence of these attractive forces, the repulsive charge is reduced and the casein micelles aggregate and eventually coagulate into a network of small chains, yoghurt coagulum, and a visco-elastic gel (Brückner-Gühmann et al., 2019).

Enrichment of yoghurt with ABP increased the viscosity of the yoghurt sample. This may have desirable effects on the eating quality of the spiced yoghurt. Spices did not have significant effects on the instrumental viscosity colour measurements. Thus, the colour attributes of the yoghurt samples were maintained. In addition, sensory evaluation revealed that the enrichment of the fermentation mix with ABP gave the highest colour, aroma, taste, texture, and overall acceptance score, as compared to that of other spices but less than that of the plain yoghurt. Colour is perceived as a measure of quality among consumers and could greatly influence the consumers' acceptability of food products. It is used to measure other quality attributes such as flavour, sensory, nutritional, and pigments, due to its simplicity and good correlation with other physicochemical properties.

Conclusions

Yoghurt containing clove extracts showed the highest antioxidant potential and a reduced rate of syneresis, but also reduced viscosity and organoleptic properties. In the case of ABP, antioxidant characteristics, protein content, and viscosity were enhanced with minimal effects on consumer acceptability. The effects of turmeric extracts include improved calcium content and reduced syneresis, but reduced total solids. Overall, African black pepper showed remarkable desirable effects on the product as its addition to the milk-based fermentation mix resulted in yoghurt which combines improved antioxidant characteristics with increased protein content and viscosity. Therefore, the enrichment of yoghurt with these food spices undoubtedly constitutes a good way for improving the nutritional value and health benefits of yoghurt. Such recipes may be considered for optimization and development towards producing new functional yoghurt.

Author Contributions: Oludare Ogunyemi, conceptualization and original draft preparation; Gideon Gyebi, methodology, writing, and reviewing; Raheemah Shaibu, data curation; Modupe Fabusiwa, writing and editing; Charles Olaiya, overall supervision

Funding: This research received no external funding

Acknowledgments: The technical support of Mr Onugwu Ernest of the Department of Biochemistry, Salem University, Nigeria and Miss Wura Subuloye of Department of Biochemistry, University of Ibadan is well acknowledged.

References

- Akalin, A., Unal, G., Dinkci, N., Hayaloglu A. (2012): Microstructural, textural, and sensory characteristics of probiotic yogurts fortified with sodium calcium caseinate or whey protein concentrate. *J. Dairy Sci.* 95, 3617-3628. <https://doi.org/10.3168/jds.2011-5297>.
- Baba, A., Najarian A., Shori A., Lit W., Keng A. (2014): Viability of Lactic Acid Bacteria, Antioxidant Activity and In Vitro Inhibition of Angiotensin-I-Converting Enzyme of *Lycium barbarum* Yogurt. *Arab. J. Sci. Eng.* 39 5355-5362. <https://doi.org/10.1007/s13369-014-1127-2>
- Beckman C. (2000): Phenolic-storing cells: keys to programmed cell death and periderm formation in wilt disease resistance and in general defence responses in plants? *Physiological and Molecular Plant Pathology* 57, 101-110. <https://doi.org/10.1006/pmpp.2000.0287>
- Brückner-Gühmann, M., Benthin, A., Drusch S. (2019): Enrichment of yoghurt with oat protein fractions:

Structure formation, textural properties and sensory evaluation. *Food Hydrocolloid.* 86, 146-153.

Caleja, C., Barros, L., Antonio, A., Carocho, M., Oliveira, M., Ferreira, I. (2016): Fortification of yogurts with different antioxidant preservatives: A comparative study between natural and synthetic additives. *Food Chem.* 210, 262-268. <https://doi.org/10.1016/j.foodchem.2016.04.114>

Cherrak, S., Mokhtari-Soulimane, N., Berroukeche, F., Bensenane, B., Cherbonnel, A., Merzouk, H., Elhabiri M. (2016): In Vitro Antioxidant versus Metal Ion Chelating Properties of Flavonoids: A Structure-Activity Investigation. *PLOS ONE* 11, e0165575-e0165575. <https://doi.org/10.1371/journal.pone.0165575>

Citta, A., Folda, A., Scalcon, V., Scutari, G., Bindoli, A., Bellamio, M., Feller, E., Rigobello M. (2017): Oxidative changes in lipids, proteins, and antioxidants in yogurt during the shelf life. *Food Sci. Nutr.* 5, 1079-1087. doi: 10.1002/fsn3.493

Corrieu G., Béal, C. (2016): Yogurt: The Product and its Manufacture. *The Encyclopedia of Food and Health* 5, 617-624. <https://doi.org/10.1016/B978-0-12-384947-2.00766-2>

Dabija, A., Codina, G., Anca, G., Sanduleac, E., Lacramioara R. (2018): Effects of some vegetable proteins addition on yogurt quality. *Scientific Study and Research: Chemistry and Chemical Engineering, Biotechnology, Food Industry* 19, 181-192.

Dusan, M., Vesna, K. (2007): Investigation of metal-flavonoid chelates and the determination of flavonoids via metal-flavonoid complexing reactions. *J. Serb. Chem. Soc.* 72, 921-939. <https://doi.org/10.2298/JSC0710921M>

El-Said M., Haggag H., Fakhr El-Din H., Gad A., Farahat A. (2014): Antioxidant activities and physical properties of stirred yoghurt fortified with pomegranate peel extracts. *Annals Agricult.Sci.* 59, 207-212. <http://dx.doi.org/10.1016/j.aoas.2014.11.007>

Gemedi, H., Haki, G., Beyene, F., Woldegiorgis, A., Rakshit S. (2015): Proximate, mineral, and antinutrient compositions of indigenous Okra (*Abelmoschus esculentus*) pod accessions: implications for mineral bioavailability. *Food Sci. Nutr.* 4, 223-233. <https://doi.org/10.1002/fsn3.282>

Güner, A., Ardic, M., Keles, A., Doğruler Y. (2007): Production of yogurt ice cream at different acidity. *Int. J. Food Sci. Technol.* 42, 948-952. <https://doi.org/10.1111/j.1365-2621.2006.01315.x>

Guler A, Mutlu B (2005): The effect of different incubation, temperature on bacteria counts of bio-yoghurt made from ewe's milk. *Int. J. Dairy Technol.* 58, 1784-1789. <https://doi.org/10.1111/j.1471-0307.2005.00209.x>

Gyebi, G., Ocheje, J., Ogunyemi O. (2019): Molecular docking studies on bioactive compounds from clove (*Syzygium Aromaticum*) on metabolic regulators in cancer. *Salem University Journal of Life Sciences* 1, 1-18.

Hashemi Gahruei, H., Eskandari, M., Mesbahi, G., Hanifpour M. (2015): Scientific and technical aspects of yogurt fortification: A review. *Food Sci. Human Wellness* 4, 1-8. <https://doi.org/10.1016/j.fshw.2015.03.002>

Jaworska, D., Rak, C., Kolanowski, W., Swiderski F. (2005): Relative importance of texture properties in the sensory quality and acceptance of natural yoghurts. *Int. J. Dairy Technol.* 58, 39-46. <https://doi.org/10.1111/j.1471-0307.2005.00178.x>

Johnson E. (1964): The Quantitative Analysis of Drugs. By D.C. Garratt. *J. Pharm. Pharmacol.* 16, 772-772. <https://doi.org/10.1111/j.2042-7158.1964.tb07408.x>

Kapadiya, D., Dabhi, B., Aparnathi, K. (2016): Spices and Herbs as a Source of Natural Antioxidants for Food. *Int. J. Curr. Microbiol. Appl. Sci.* 5, 280-288. <https://doi.org/10.20546/ijcmas.2016.507.029>

Khan, I., Bule, M., Ullah, R., Nadeem, M., Asif, S., Niaz, K. (2019): The antioxidant components of milk and their role in processing, ripening, and storage: Functional food. *Vet. World* 12, 12-33. <https://doi.org/10.14202/vetworld.2019.12-33>

Latimer, G. Association of Official Analytical Chemists I. (2019): *Official methods of analysis of AOAC International*, Gaithersburg, Maryland: AOAC International.

Lee, W., Lucey, J. (2010): Formation and Physical Properties of Yogurt. *Asian-Australia. J. Anim. Sci.* 23, 1127 - 1136. <https://doi.org/10.5713/ajas.2010.r.05>

Morell, P., Hernando, I., Llorca, E., Fiszman S. (2015): Yoghurts with an increased protein content and physically modified starch: Rheological, structural, oral digestion and sensory properties related to enhanced satiating capacity. *Food Res. Int.* 70, 64-73. <https://doi.org/10.1016/j.foodres.2015.01.024>

Muniandy, P., Shori, A., Baba, A. (2016): Influence of green, white and black tea addition on the antioxidant activity of probiotic yogurt during refrigerated storage. *Food Pack. Shelf Life* 8, 1-8. <https://doi.org/10.1016/j.fpsl.2016.02.002>

Najgebauer-Lejko, D., Sady, M. (2015): Estimation of the antioxidant activity of the commercially available fermented milks. *Acta Sci. Pol. Technol.* 14, 387-396. <https://doi.org/10.17306/J.AFS.2015.4.38>

Ndife, J. (2014): Production and Quality Assessment of Functional Yoghurt Enriched with Coconut. *Int. J. Food Sci. Nutr.* 3, 545-550. <https://doi.org/10.11648/j.ijfnfs.20140306.19>

Ogunyemi, O., Gyebi, G., Shaibu, O., Ozonwa, O., Olaiya C. (2020): Enhancement of antioxidant potential of yoghurt with natural additives: preliminary sensory and in vitro antioxidant analyses. *Salem University Journal of Life Sciences* 2, 1-17.

Olaiya, C., Choudhary, M., Ogunyemi, O., Nwauzoma, A. (2013): Nutraceuticals from Bitter Leaf (*Vernonia amygdalina* Del.) Protects against Cadmium Chloride induced Hypertension in Albino Rats. *Nat. Sci.* 11, 136-145.

Oyaizu, M. (1986): Studies on products of browning reaction antioxidative activities of products of

browning reaction prepared from glucosamine. *Jpn. J. Nutr.* 44, 307-316.

Panche, A., Diwan, A., Chandra S. (2016): Flavonoids: an overview. *J. Nutr. Sci.* 5, e47-e47. 10.1017/jns.2016.41

Raikos, V., Ni, H., Hayes, H., Ranawana V. (2018): Antioxidant Properties of a Yogurt Beverage Enriched with Salal (*Gaultheria shallon*) Berries and Blackcurrant (*Ribes nigrum*) Pomace during Cold Storage. *Beverages* 5(1), 2. <https://doi.org/10.3390/beverages5010002>

Ramaswamy, H., Basak S. (1992): Pectin and Raspberry Concentrate Effects on the Rheology of Stirred Commercial Yogurt. *J. Food Sci.* 57, 357-360. <https://doi.org/10.1111/j.1365-2621.1992.tb05494.x>

Re, R., Pellegrini, N., Proteggente, A., Pannala, A., Yang, M., Rice-Evans C. (1999): Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radic. Biol. Med.* 26, 1231-1237.

Ruberto, G., Baratta, M., Deans, S., Dorman, H. (2000) Antioxidant and antimicrobial activity of *Foeniculum vulgare* and *Crithmum maritimum* essential oils. *Planta Med.* 66, 687-693. <https://doi.org/10.1055/s-2000-9773>

Senadeera, S., Prasanna, P., Jayawardana, N., Gunasekara, D., Senadeera, P., Chandrasekara A. (2018): Antioxidant, physicochemical, microbiological, and sensory properties of probiotic yoghurt incorporated with various *Annona* species pulp. *Helijon* 4, e00955. <https://doi.org/10.1016/j.heliyon.2018.e00955>

Shirwaikar, A., Rajendran, K., Punitha I. (2006): In vitro antioxidant studies on the benzyl tetra isoquinoline alkaloid berberine. *Biol. Pharm. Bull.* 29, 1906-1910.

Singleton, V., Orthofer, R., Lamuela-Raventós, R. (1999) Analysis of total phenols and other oxidation substrates and antioxidants by means of folin-ciocalteu reagent. *Meth. Enzymol.* 299, 152-178. [https://doi.org/10.1016/S0076-6879\(99\)99017-1](https://doi.org/10.1016/S0076-6879(99)99017-1)

Soetan, K., Olaiya, C., Oyewole, O. (2009): The importance of mineral elements for humans, domestic animals and plants: A review. *Afr. J. Food Sci.* 4, 200-222.

Srivastava, P., Prasad, S., Ali, M., Prasad, M. (2015): Analysis of antioxidant activity of herbal yoghurt prepared from different milk. *The Pharma Innovation Journal* 4, 18-20.

Tamang, J., Shin, D., Jung, S., Chae S. (2016): Functional Properties of Microorganisms in Fermented Foods. *Front. Microbiol.* 7, 578-578. <https://doi.org/10.3389/fmicb.2016.00578>

Tavakoli, M., Habibi Najafi, M., Mohebbi, M. (2019): Effect of the milk fat content and starter culture selection on proteolysis and antioxidant activity of probiotic yogurt. *Helijon* 5, e01204-e01204. <https://doi.org/10.1016/j.heliyon.2019.e01204>

Tomovska, J., Gjorgievski N., Makarijoski, B. (2016). Examination of pH, Titratable Acidity and Antioxidant Activity in Fermented Milk. *J. Mater. Sci. Eng.* 6, 326-333. <https://doi.org/10.17265/2161-6213/2016.11-12.006>

Wibawanti, M., Rinawidiastuti, Arifin, D., Zulfanita. (2018) Improving characteristics of goat milk yogurt drink fortified by mangosteen rind (*Garcinia mangostana* Lin .) extract. *IOP Conference Series: Earth and Environmental Science* 102, <https://doi.org/10.1088/1755-1315/102/1/012008>.

Zhishen, J., Mengcheng, T., Jianming, W. (1999) The determination of flavonoid contents in mulberry and their scavenging effects on superoxide radicals. *Food Chem.* 64, 555-559. [https://doi.org/10.1016/S0308-8146\(98\)00102-2](https://doi.org/10.1016/S0308-8146(98)00102-2)