



## Feasibility study of a natural vinaigrette from olive (*Olea europaea*) oil and lemon (*Citrus limonium*) juice-based lemon albedo extract

Malika Chekroune and Salem Benamara\*

University M'hamed Bougara, Faculty of Engineering Sciences, Department of Process Engineering, Research Laboratory in Food Technology, Avenue de l'Indépendance, Boumerdès, 35000, Algeria

### ARTICLE INFO

#### Article history:

Received: August 21, 2020

Accepted: May 4, 2021

#### Keywords:

albedo

emulsion

lemon (*Citrus limonium*)

olive oil

vinaigrette stability

### ABSTRACT

The present paper reports on the feasibility study of a natural vinaigrette from olive (*Olea europaea*) oil and lemon (*Citrus limonium*) juice-based lemon albedo extract. The albedo extract was prepared at different albedo to lemon juice ratios. The physical stability of different vinaigrette samples, in terms of creaming behaviour, was adopted as quality criterion. The most stable sample was then analysed for its rheological behaviour. According to preliminary results, the most physically stable vinaigrette could be obtained at albedo: lemon juice ratio of 1:15.1 (w/w). The corresponding dressing showed a stability of six days and a shear-thinning type rheological behaviour. So, the formulation of natural vinaigrette from olive oil, lemon juice and lemon albedo is feasible but such formulation is of short-term physical stability, which is a general well-known characteristic of natural and minimally processed food products. In our opinion, this study merits further research, in order to: i) better optimize the vinaigrette formulation, and ii) characterize the functional properties of the final product.

### Introduction

In recent years, nutrition has been considered as a determining factor in sustaining and restoring health. Various processed food products have been incriminated in the appearance of severe diseases such as cancer, hypertension, diabetes, etc. (Medeiros et al., 2012). In particular, Ngo et al. (2011) brought up the undesirable effects of synthetic ingredients on human health.

Emulsifiers represent a group of essential ingredients used to stabilize heterogeneous liquid systems. In the case of olive-in-water emulsions, emulsifiers are usually added to the aqueous phase to improve their stability against creaming.

Polysaccharides are known to be among the most employed emulsifiers in food industry. Pectins, sub-group molecules belonging to polysaccharides, are widespread in citrus peels. They have many functional properties such as the stabilizing power of food emulsions. These molecules are obtained

in purified form via numerous technological operations implicating various chemical extraction agents (Seixas et al., 2014). On the other hand, there is an emphasis on investigating natural emulsifiers that can be efficiently incorporated into various food emulsions (Yang & McClements, 2013). As pointed out by Rezzadori et al. (2012) about pectin extraction, the idea is to apply an alternative process that does not generate toxic waste. In this context, Fidalgo et al. (2016) have studied the simultaneous extraction of pectin and essential oils from orange and lemon peels, using only water as dispersing medium. Banerjee et al. (2016) have evaluated the effect of lemon juice on pectin extraction from mango peels.

While the instantly prepared lemon olive oil vinaigrette is now very popular, particularly in the Mediterranean countries, there is, to our knowledge, no work about the stabilisation of such products with only natural raw ingredients, including emulsifiers. In this respect, the few studies devoted to this subject have used purified polysaccharides as stabilizers (Abedinzadeh et al.,

\*Corresponding author E-mail: [sbenamara2001@yahoo.fr](mailto:sbenamara2001@yahoo.fr)

2016; Silva et al., 2010; Paraskevopoulou et al., 2007; Paraskevopoulou et al., 2005).

The present paper reports on the feasibility study for natural vinaigrette from olive (*Olea europaea*) oil and lemon (*Citrus limonium*) juice-based lemon albedo extract (LAE), at olive oil:LAE ratio of 1:1.2 (v/v). Such natural formulations, despite their short shelflife, refer currently to a major trend at global level (Silveira et al., 2013).

## Materials and methods

### Materials

The virgin olive oil from *Chemlal* olive variety was collected in Bouira region (120 km south east of Algiers).

Lemon albedo was recovered by peeling local lemon fruits. The albedo was separated from coloured peel (flavedo), cut in small size pieces (~2mm) and then dried at 50 °C until a constant weight.

The lemon juice was freshly extracted using a home lemon-press and then centrifuged.

### Vinaigrette Preparation

In total, 10 vinaigrette samples were prepared, according to diagram in Figure 1, proposed here for the first time to our knowledge.

The LAE was prepared by mixing 145 mL hot lemon juice with lemon albedo small pieces at the following mixing ratios (w/v): 1:24.2, 1:20.7, 1:18.1, 1:16.1, 1:15.6, 1:15.1, 1:14.6, 1:14.1, 1:13.7 and 1:13. The homogenization was performed using an Ultra-Turrax T25 homogenizer (IKA Instrument, Germany) equipped with a dispersing tool (S25-N). Shortly after, each vinaigrette sample was divided equally into six 15 mL- polypropylene tubes with a conical bottom which were then stored at room temperature to monitor the physical stability.

### Physical Stability Analysis

As underlined by Sosa et al. (2014), the stability of food emulsions is a key factor in the development of such products. The physical stability in particular indicates the ability of emulsion to resist changes in spatial distribution of ingredients over time (McClements, n.d), thus mainly determining the shelf life of the product (Junqueira et al., 2018).

The stability was estimated by monitoring over time of the creaming behaviour (demixing kinetics). For this, the serum volume separated on the bottom of tubes was measured at different time intervals and the curve fitting was performed using Excel software.

The lower the creaming rate, the more stable the vinaigrette.

The most stable vinaigrette was further analysed for its rheological behaviour. A constant rheological behaviour over time indicates a better physical stability.

The rheological analysis was performed using the VT550 viscotester (Thermo Haake). The flow curves of vinaigrette samples were obtained at 20 °C at various shear rates (0-400 s<sup>-1</sup>) for five minutes.


## Results and discussion

The creaming behaviour of obtained vinaigrette samples is illustrated in Figure 2a. As can be seen, straight lines are obtained for all samples, except for those with a lemon albedo:lemon juice ratio below 1:20.7 (w/v), indicating the influence of albedo proportion on the creaming kinetic. We think that the pectin could be the determining factor in the emulsifying ability of the lemon albedo. It is worthwhile to note that the citrus pectin, as a natural hydrocolloid (Korish, 2015) could be an interesting alternative emulsifier, stabilizer and gelling agent (Gull et al., 2018), considering its capacity, as hydrocolloid, to increase the viscosity of multiphase systems (Venir and Maltini, 2013, Prabhuswamy et al., 2019). Our findings are in concordance with those communicated by Santiago et al. (2002) regarding the effect of polysaccharide concentration on the creaming kinetics of oil-in-water emulsion.

The best stability that has been reached is about six days and corresponds to the creaming rate very close to zero. It was obtained with the albedo:lemon juice mixing ratio of 1:15.1 (w/v) that is 9.6 g lemon albedo for 145 mL boiled lemon juice. This stability is close to that (four days) communicated by Perrechil et al. (2014) about an emulsion stabilized by sodium caseinate- locust bean gum Maillard conjugates used as substitutes of synthetic emulsifiers. Despite the short shelf life of natural and minimally processed products, many researchers encourage their consumption regarding their positive effects on health (De Podesta et al., 2007).

It should be noted that in our case, the lemon albedo:lemon juice ratio in the LAE cannot be increased beyond 1:15.1 (w/v), otherwise the vinaigrette loses the fluidity characteristic of this type of product. At last, a full understanding of the emulsifying-stabilizing effect of the LAE is not so easy, knowing the structural complexity of the oil-in-water emulsions (Johnston et al., 2015).


The flow curves of the most stable vinaigrette, at different storage times, are presented in Figure 2b.



**Figure 1.** Diagram of preparation of natural vinaigrette samples

From Figure 1, the selected formula seems to safeguard its rheological stability at shear rates not exceeding 40 (1/s). At this shear rate, the overall shape of flow curves does not change during the storage period of six days. As highlighted by Bosco et al. (2019), regarding rheological parameters of shear-thickening fluids, this observation may be linked to the fact that low shear rates marginally impact the viscosity values and thus the

microstructure of vinaigrette. The regression of  $\ln(\text{shear stress } (\tau))$  versus  $\ln(\text{shear rate } (\dot{\gamma}))$  (not presented here), allowed us to deduce the values (below 1) of behaviour index  $n$ , demonstrating a rheological behavior of shear-thinning type. These findings are consistent with those reported by Nikovska (2012) concerning olive oil-in-water emulsions with lower protein emulsifier concentration.



**Figure 2.** Serum Volume of separated serum versus storage time, for all tested vinaigrette samples (a), and flow curves at different storage times of the most stable vinaigrette sample (b).

## Conclusions

Results show that the formulation of natural vinaigrette from olive oil, lemon juice and lemon juice-based lemon albedo extract is feasible, by choosing the appropriate proportions of components. The best physical stability reached is of six days, knowing that the short-term stability is a general characteristic of natural processed food products.

In our opinion, this study needs to be developed, in order to: i) better optimize the vinaigrette formulation, and ii) characterize the functional properties of the final product.

**Author Contributions:** Chekroune M. conceived, designed and performed the experiments. Benamara S. supervised the work and he rewrote the manuscript by updating the bibliographic research.

**Funding:** This research received no external funding

## References

- Abedinzadeh, S., Torbati, M., Azadmard-Damirchi, S. (2016): Some Qualitative and Rheological Properties of Virgin Olive OilApple Vinegar Salad Dressing Stabilized With Xanthan Gum. *Adv. Pharm. Bull.* 6 (4), 597-606. <https://doi.org/10.15171%2Fapb.2016.074>
- Banerjee, J., Vijayaraghavan, R., Arora, A., MacFarlane, D. R., Patti, A. F. (2016): Lemon juice based extraction of pectin from mango peels: waste to wealth by sustainable approaches. *ACS Sustain. Chem. Eng.* 4 (11), 5915-5920. <https://doi.org/10.1021/acssuschemeng.6b01342>
- Bosco, A., Calado, V., Maia, J. (2019): Rheological parameters of shear-thickening fluids using an experimental design. *Mater. Res.* 22 (5): e20180631. <https://doi.org/10.1590/1980-5373-MR-2018-0631>
- De Podesta, O.P.G, Peres, S.V., Salaroli, L.B., Cattafesta, M., De Podesta, J.R.V., von Zeidler, S.L.V., de Oliveira, J.C., Kowalski, L.P., Ikeda, M.K., Brennan, P., Curado, M.P. (2019): Consumption of minimally processed foods as protective factors in the genesis of squamous cell carcinoma of the head and neck in Brazil. *PLoS One* 14 (7), e0220067. <https://doi.org/10.1371/journal.pone.0220067>

- Fidalgo, A., Ciriminna, R., Carnaroglio, D., Tamburino, A., Cravotto, G., Grillo, G., Ilharco, L. M., Pagliaro, M. (2016): Eco-friendly extraction of pectin and essential oils from orange and lemon peels. *ACS Sustain. Chem. Eng.* 4 (4), 2243-2251. <https://doi.org/10.1021/acssuschemeng.5b01716>
- Gull, A., Kumar, P., Safapuri, T. A., Khajuria, A. K., Gandotra, A., Jan, R. (2018): The effect of adding non-conventional ingredients and hydrocolloids to desirable quality attributes of pasta. A Mini Review. *Croat. J. Food Sci. Technol.* 10 (1), 106-111. <https://doi.org/10.17508/CJFST.2018.10.1.08>
- Johnston, S. P., Nickerson, M. T., Low, N. H. (2015): The physicochemical properties of legume protein isolates and their ability to stabilize oil-in-water emulsions with and without genipin. *J. Food Sci. Technol.* 52, 4135-4145. <https://doi.org/10.1007/s13197-014-1523-3>
- Junqueira, L. A., Amaral, T. N., Oliveira, N. L., Prado, M. E. T., de Resende, J. V. (2018): Rheological behavior and stability of emulsions obtained from *Pereskia aculeata* Miller via different drying methods. *Int. J. Food Prop.* 21 (1), 21-35. <https://doi.org/10.1080/10942912.2018.1437177>
- Korish, M. (2015): Potential utilization of *Citrullus lanatus* var. *Colocynthoides* waste as a novel source of pectin. *J. Food Sci. Technol.* 52 (4), 2401-2407. <https://doi.org/10.1007/s13197-014-1277-y>
- McClements, D. J. (n.d): Stability of Food Emulsions (1). [https://people.umass.edu/~mcclemen/FoodEmulsions2008/Presentations\(PDF\)/\(6\)Emulsion\\_Stability.pdf](https://people.umass.edu/~mcclemen/FoodEmulsions2008/Presentations(PDF)/(6)Emulsion_Stability.pdf). Accessed August 11, 2020.
- Medeiros, R. A., Lourencao, B. C., Rocha-Filho, R. C., Fatibello-Filho, O. (2012): Simultaneous voltammetric determination of synthetic colorants in food using a cathodically pretreated boron-doped diamond electrode. *Talanta* 97 (15), 291-297. <https://doi.org/10.1016/j.talanta.2012.04.033>
- Ngo, D.-H., Wijesekara, I., Vo, T. S., Ta, Q. V., Kim, S.-K. (2011): Marine food-derived functional ingredients as potential antioxidants in the food industry: An overview. *Food Res. Int.* 44 (2), 523-529. <https://doi.org/10.1016/j.foodres.2010.12.030>
- Nikovska, K. (2012): Study of olive oil-in-water emulsions with protein emulsifiers. *Emir. J. Food Agric.* 24 (1), 17-24. <https://doi.org/10.9755/ejfa.v24i1.10594>
- Paraskevopoulou, A., Boskou, D., Kiosseoglou, V. (2005): Stabilization of olive oil – lemon juice emulsion with polysaccharides. *Food Chem.* 90 (4), 627-634. <https://doi.org/10.1016/j.foodchem.2004.04.023>
- Paraskevopoulou, D., Boskou, D., Paraskevopoulou, A. (2007): Oxidative stability of olive oil–lemon juice salad dressings stabilized with polysaccharides. *Food Chem.* 101 (3), 1197-1204. <https://doi.org/10.1016/j.foodchem.2006.03.022>
- Perrechil, F. A., Santana, R.C., Lima, D. B., Polastro, M. Z., Cunha, R. L. (2014): Emulsifying properties of maillard conjugates produced from sodium caseinate and locust bean gum. *Braz. J. Chem. Eng.* 31 (2), 429-438. <https://doi.org/10.1590/0104-6632.20140312s00002328>
- Prabhuswamy, T., Tamilselvan, K., Michael, M.L., Moses, J.A., Chinnasamy, A. (2019): Influence of different hydrocolloids on dispersion of sweet basil seeds (*Ocimum basilicum*) in fruit-flavoured beverages. *Croat. J. Food Sci. Technol.* 11 (1), 37-43. <https://doi.org/10.17508/CJFST.2019.11.1.05>
- Rezzadori, K., Benedetti, S., Amante, E.R. (2012): Proposals for the residues recovery: Orange waste as raw material for new products. *Food Bioprod. Process.* 90 (4), 606-614. <https://doi.org/10.1016/j.fbp.2012.06.002>
- Santiago, L.G., Gonzalez, R.J., Fillery-Travis, A., Robins, M., Bonaldo, A.G., Carrara, C. (2002): The influence of xanthan and  $\lambda$ -carrageenan on the creaming and flocculation of an oil-in-water emulsion containing soy protein. *Braz. J. Chem. Eng.* 19 (4), 411-417. <https://doi.org/10.1590/S0104-66322002000400009>
- Seixas, F.L., Fukuda, D.L., Turbiani, F.R.B., Garcia, P.S.; Petkowicz, C.L. de O., Jagadevan, S., Gimenes, M. L. (2014): Extraction of pectin from passion fruit peel (*Passiflora edulis* f. *flavicarpa*) by microwave-induced heating. *Food Hydrocoll.* 38, 186-192. <https://doi.org/10.1016/j.foodhyd.2013.12.001>
- Silva, K.A., Rocha-Leão, M.H., Coelho, M.A. Z. (2010): Evaluation of aging mechanisms of olive oil-lemon juice emulsion through digital image analysis. *J. Food Eng.* 97 (3), 335-340. <https://doi.org/10.1016/j.jfoodeng.2009.10.026>
- Silveira, A.C., Aguayo, E., Artés, F. (2013): Shelf-life and quality attributes in fresh-cut Galia melon combined with fruit Juices. *LWT - Food Sci. Technol.* 50 (1), 343-348. <https://doi.org/10.1016/j.lwt.2012.04.010>
- Sosa, N., Schebor, C., Pérez, O.E. (2014): Encapsulation of citral in formulations containing sucrose or trehalose: Emulsions properties and stability. *Food Bioprod. Process.* 92, 266-274. <https://doi.org/10.1016/j.fbp.2013.08.001>
- Venir, E., Maltini, E. (2013): Relevance of physical properties in the stability of plant-food based products. *Indian J. Exp. Biol.* 51, 895-904.
- Yang, Y., McClements, D.J. (2013): Encapsulation of vitamin E in edible emulsions fabricated using a natural Surfactant. *Food Hydrocoll.* 30 (2), 712-720. <https://doi.org/10.1016/j.foodhyd.2012.09.003>